



Year: 2017

A cross-sectional observational study of high override rates of drug allergy alerts in inpatient and outpatient settings, and opportunities for improvement

Slight, Sarah Patricia ; Beeler, Patrick E ; Seger, Diane L ; Amato, Mary G ; Her, Qoua L ; Swerdloff, Michael ; Dalleur, Olivia ; Nanji, Karen C ; Cho, InSook ; Maniam, Nivethietha ; Eguale, Tewodros ; Fiskio, Julie M ; Dykes, Patricia C ; Bates, David W

Abstract: **OBJECTIVES:** To evaluate how often and why providers overrode drug allergy alerts in both the inpatient and outpatient settings. **DESIGN:** A cross-sectional observational study of drug allergy alerts generated over a 3-year period between 1 January 2009 and 31 December 2011. **SETTING:** A 793-bed tertiary care teaching affiliate of Harvard Medical School and 36 primary care practices. **PARTICIPANTS:** Drug allergy alerts were displayed for a total of 29 420 patients across both settings. **MAIN OUTCOME MEASURES:** Proportion of drug allergy alerts displayed and overridden, proportion of appropriate overrides, proportion of overrides in each medication class, different reasons for overriding and types of reactions overridden. **RESULTS:** A total of 158 023 drug allergy alerts were displayed, 131 615 (83%) in the inpatient setting and 26 408 (17%) in the outpatient setting; 128 157 (81%) of which were overridden. A random sample of inpatient (n=200, 0.19%) and outpatient (n=50, 0.25%) alert overrides were screened for appropriateness, with >96% considered appropriate. Alerts for some drug classes, such as 'non-antibiotic sulfonamides', were overridden for >81% of prescriptions in both settings. The most common override reason was patient has taken previously without allergic reaction. In the inpatient setting alone, 70.9% of alerts that warned against the risk of anaphylaxis were overridden. **CONCLUSIONS:** The information contained in patients' drug allergy lists needs to be regularly updated. Most of the drug allergy alerts were overridden, with the majority of alert overrides in the subsample considered appropriate. Some of the rules for these alerts should be carefully reviewed and modified, or removed. Further research is needed to understand providers' overriding of alerts that warned against the risk of 'anaphylaxis', which are more concerning with respect to patient safety.

DOI: <https://doi.org/10.1136/bmjqs-2015-004851>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-123958>

Journal Article

Accepted Version

Originally published at:

Slight, Sarah Patricia; Beeler, Patrick E; Seger, Diane L; Amato, Mary G; Her, Qoua L; Swerdloff, Michael; Dalleur, Olivia; Nanji, Karen C; Cho, InSook; Maniam, Nivethietha; Eguale, Tewodros; Fiskio, Julie M; Dykes, Patricia C; Bates, David W (2017). A cross-sectional observational study of high override rates of drug allergy alerts in inpatient and outpatient settings, and opportunities for improvement. *BMJ Quality and Safety*, 26(3):217-225.

DOI: <https://doi.org/10.1136/bmjqs-2015-004851>

A Cross-Sectional, Observational Study Of High Override Rates Of Drug Allergy Alerts In Inpatient And Outpatient Settings and Opportunities for Improvement

ABSTRACT

Objectives: To evaluate how often and why providers overrode drug allergy alerts in both the inpatient and outpatient settings.

Design: A cross-sectional, observational study of drug allergy alerts generated over a three-year period between Jan 1st, 2009, and Dec 31st, 2011.

Setting: A 793-bed tertiary care teaching affiliate of Harvard Medical School and 36 primary care practices.

Participants: Drug allergy alerts were displayed for a total of 29,420 patients across both settings.

Main outcome measures: Proportion of drug allergy alerts displayed and overridden, proportion of appropriate overrides, proportion of overrides in each medication class, different reasons for overriding, and types of reactions overridden.

Results: A total of 158,023 drug allergy alerts were displayed, 131,615 (83%) in the inpatient setting and 26,408 (17%) in the outpatient setting; 128,157 (81%) of which were overridden. A random sample of inpatient (n=200, 0.19%) and outpatient (n=50, 0.25%) alert overrides were screened for appropriateness, with over 94% considered appropriate. Some drug classes, such as 'Non-antibiotic sulfonamides', were overridden over 81% of the time in both settings. The most common override reason was '*patient has taken previously without allergic reaction*'. In the inpatient setting alone, 70.9% of alerts that warned against the risk of '*anaphylaxis*' were overridden.

Conclusions: The information contained in patients' drug allergy lists needs to be regularly updated. Most of the drug allergy alerts were overridden, with the majority of alert overrides in the sub-sample considered appropriate. Some of the rules for these alerts should be carefully reviewed and modified, or removed. Further research is needed to understand providers' overriding of alerts that warned against the risk of 'anaphylaxis', which are more concerning with respect to patient safety.

INTRODUCTION

The prevalence of allergies is increasing not only in the U.K. but across the world, with as many as 30% of adults and 40% of children affected by allergies in the United States (U.S.).¹ Drug allergies in particular are becoming increasingly common. It has been estimated that penicillin in particular is responsible for approximately 75% of fatal anaphylactic cases in the U.S., leading to 500 - 1,000 deaths per year.²

Computerized physician order entry (CPOE) with clinical decision support (CDS) allows physicians to enter medication orders electronically and provides real time guidance and support to assist them with their prescribing.³⁻⁶ This guidance can take many different forms, including visual alerts or reminders about potential hazardous drug-drug or drug allergy interactions.^{7,8} However, the problem of alert-fatigue has been well documented; prescribers are often exposed to too many alerts, which can result in them ignoring or failing to respond appropriately to even important alerts which can result in serious harm.^{9,10}

Drug allergy alerts differ from other types of medication-related alerts in that they are heavily dependent on patient recall rather than those that depend on discrete lab values, diagnoses, or known interactions with other drugs.¹¹ In CPOE systems, prescribed drugs can be checked against the patient's allergy list, and a drug allergy alert can warn the prescriber about a possible 'drug allergy'. A 'drug allergy' is a type of unpredictable drug hypersensitivity reaction, which can be either immediate (immunoglobulin E (IgE)-mediated) or delayed (non-IgE-mediated).¹² These allergic reactions may often present with a broad range of symptoms, including prominent physical signs such as an itchy red skin rash (e.g., urticarial), nausea, vomiting or mental status changes. Drug-induced anaphylaxis is potentially fatal and usually

characterized by rapid-onset cardiovascular collapse. A number of studies have reported very high override rates of drug allergy alerts in primary care.^{9,13} However, it is currently unclear which drug classes and types of reactions were highly overridden. Another study recently evaluated providers' drug allergy alert overrides in two U.S. hospitals.¹⁴ However, this study does not consider the appropriateness of providers' drug allergy alert overrides, or how often these specific drug classes are overridden in primary care.

Another factor complicating safety is the existence of cross-sensitization. Once an individual has become sensitized to a drug, the possibility exists that the same individual may also react to drugs with a close structural chemical relationship or to immunochemically similar metabolites.¹² Drug allergy alerts may be generated for *possible* cross-reactivity associations between drugs, such as penicillins and other antibiotics containing a beta-lactam ring e.g., cephalosporins, carbapenems, and monobactams.¹⁵ According to the World Allergic Organization, the range of cross-sensitization varies greatly among individuals and is often difficult to predict with any certainty.¹² In this study, we evaluated how often and more importantly why providers overrode drug allergy alerts in both the inpatient and outpatient settings. Specific outcomes were: (1) the appropriateness of providers' drug allergy alert overrides, (2) proportion of overrides in each medication class, (3) the reasons why providers chose to override these alerts, and (4) the different types of reactions overridden.

METHODS

Research study site

This study included data from the Brigham and Women's Hospital (Boston,

MA), a 793-bed tertiary care teaching affiliate of Harvard Medical School, and 36 primary care practices. All these health organizations are part of a regional integrated healthcare delivery system, Partners HealthCare.

Electronic Health Record (EHR) and Clinical Decision Support

At the time of this study, all prescriber orders were entered through the Brigham Integrated Clinical Information System (BICS) at the Brigham and Women's hospital, and the Longitudinal Medical Record (LMR) in the ambulatory care setting. The BICS and LMR provide clinical, administrative, and financial computing functions, which included patient-specific decision support usually presented to providers in the form of medication alerts at the time of prescribing. These include drug-allergy interaction alerts, which are triggered if the patient has a documented allergy or intolerance in their EHR to the prescribed product or drug agent. Drug allergy alert logic was initially sourced from the commercial knowledge base, First DataBank (First DataBank, Inc. South San Francisco, CA), and reviewed by an expert committee. In contrast to the Partners drug-drug interaction knowledge base, the drug allergy alert knowledge base was not been modified or iteratively improved over time to reduce the number of false positive alerts.

When a drug allergy interaction alert is generated in either the inpatient or outpatient setting, a specific recommendation is presented to the physician, which is linked to a monograph. The alert message will indicate whether the patient has a '*definite*' (the drug being ordered is an exact match to the allergen e.g., penicillin documented in allergy list and penicillin ordered), '*probable*' (the drug being ordered is in the same allergen group e.g., penicillin (penicillins) documented in allergy list and amoxicillin (penicillins) ordered), or '*possible*' reaction (cross sensitivity is

considered likely e.g., penicillin (penicillins) documented in allergy list and cephalexin (cephalosporins) ordered), as well as the specific type of reaction that the patient may be at risk of developing e.g., hives, anaphylaxis (Figures 1 and 2).

Drug-Allergy Alerts

YOU ARE WRITING FOR:
CODEINE SULFATE TABS PO

Cancel Current Order

To keep current order, you must respond to the interaction alerts below.

Drug-Allergy Interaction Alerts

Pt. has a PROBABLE allergy to Percocet (OXYCODONE HCL/ACETAMINOPHEN); reaction is Anaphylaxis.

Pt. has a DEFINITE allergy to codeine; reactions are Anaphylaxis, Rash, Flushing.

Reason for override:

Previous override:

Keep Current Order

Figure 1. Screenshot of a drug allergy alert in the inpatient setting

Warning

You are ordering: CODEINE SULFATE TABS

Drug - Allergy Intervention

Alert Message: The patient has a definite allergy: codeine. Reaction(s): Anaphylaxis.

Keep New Order - select reason(s)

☒ Patient does not have this allergy, will D/C pre-existing allergy

Reasons for override:

☐ Patient has taken previously without allergic reaction

☐ Low risk cross sensitivity, will monitor

☐ No reasonable alternatives

☐ Other

Continue Order Cancel Back To Search

Figure 2. Screenshot of a drug allergy alert in the outpatient setting

In the outpatient setting, the alert warning gives the provider the choice of selecting '*patient does not have this allergy, will discontinue pre-existing allergy*' or overriding the alert. Should the physician choose to override the alert, they are

required to select one of the following coded ‘override’ reasons in order to proceed: *‘Patient has taken previously without allergic reaction’, ‘Low risk cross sensitivity, will monitor’, ‘No reasonable alternatives’,* or *‘Other’* (Figure 2). If *‘Other’* is selected, further details should be entered in the free-text field. In the inpatient setting, the alert warning gives the provider the choice of cancelling the current order or keeping it, in which case they are required to enter a reason justifying their decision to override the alert in the free-text field provided (Figure 1).

Study design and sample selection

This study was a cross-sectional, observational study of drug allergy alerts generated over a three-year period between January 1st, 2009, and December 31st, 2011. The necessary approvals were obtained to access these data from the Partners Human Research Committee (PHRC), which is the Institutional Review Board (IRB) of Partners Research Management at Partners HealthCare. All drug allergy alerts triggered in the inpatient and outpatient settings during the study period were downloaded. This included patients’ names and medical record identification numbers; names of both the allergic reaction and medicine that triggered the drug allergy alert; date of alert; practice location (outpatients)/specialty (inpatients); and the reasons given by providers at the time of overriding the alert.

Appropriateness check

Alert overrides of drug allergy alerts were considered appropriate for new orders in the inpatient and outpatient settings, if the reason indicated by the provider e.g., *‘patient does not have this allergy, will discontinue pre-existing allergy’, ‘low risk cross sensitivity, will monitor’, ‘no reasonable alternatives’, ‘patient has taken*

previously without allergic reaction, or *other* could be verified on chart review. For renewal orders, the patient's chart was reviewed to confirm that there was no previous history that the patient had an allergic reaction to the drug (e.g., rash, hives, anaphylaxis, mental status change, nausea and vomiting) and the patient did not experience an adverse event after the medication was ordered for the patient.

A pharmacist (D.L.S.) screened a random sample of inpatient (n=200, 0.19%) and outpatient (n=50, 0.25%) alert overrides for appropriateness. A second academic pharmacist (M.A.) independently reviewed the electronic inpatient medical charts (n=200), and inter-rater agreement comparing the appropriateness versus inappropriateness of each alert override calculated and found to be excellent ($\kappa=0.76$). An attending physician (K.C.N) and academic nurse (I.C.) independently reviewed the electronic outpatient medical charts (n=50), and inter-rater agreement was also found to be excellent ($k=0.86$). Any disagreements were resolved by discussion with another reviewer (D.W.B.).

Data Analysis

Data were processed using structured query language (SQL) statements. Natural language processing algorithms were developed to complete the categorizations of free-text override reasons in the outpatient setting and uncoded allergic reactions in both settings. For alerts that displayed more than one different type of allergic reactions, only the most severe reactions were considered. All drugs triggering allergy alerts were mapped to ATC codes (Anatomical Therapeutic Chemical Classification System, World Health Organization, Geneva, Switzerland) in order to enable grouping of drugs into drug classes. In the inpatient setting, a research assistant (N.M.) mapped each of the free-text entries given by providers to justify

their decision to override the alert (n=20,699) to one of 30 individual categories. Each category contained a group of words or phrases with similar meaning or connotations, and included the four coded reasons in the outpatient setting listed above for comparison. Words and phrases that could not be grouped were classified under 'Other'. D.L.S provided a second check of his classification process. Descriptive statistics were computed using the software R, version 3.0.2 (R Foundation for Statistical Computing, Vienna, Austria).

RESULTS

Summary

A total of 158,023 drug allergy alerts were displayed, 131,615 (83%) in the inpatient setting and 26,408 (17%) in the outpatient setting, over the three-year time period; 128,157 (81%) of which were overridden. These alerts were displayed for a total of 29,420 patients across both settings. The drug classes 'Opioids', 'Non-antibiotic sulfonamides (without coxibs)', which included drugs like furosemide (Lasix®) and indapamide (Lozol®), and 'Non-antibiotic sulfonamides (coxibs)', which included drugs like celecoxib (Celebrex®), were overridden over 81% of the time in both settings. The most common reason for overriding drug allergy alerts in both settings was '*patient has taken previously without allergic reaction*'. However, in a random sample of inpatient (n=200, 0.19%) and outpatient (n=50, 0.25%) alert overrides, over 94% (n=250) of alert overrides in both settings were considered appropriate. In the inpatient setting, 70.9% (n=1,682) of alerts that displayed '*anaphylaxis*' were overridden, compared to 56% (n=130) in the outpatient setting.

Number, type and overrides of drug allergy alerts

A total of 131,615 drug allergy alerts were displayed in the inpatient setting over the study time period. The provider indicated that the '*patient did not have this allergy*' for 2% (n=2,602) of these alerts. The overall alert override rate was 81.9% (n=107,812) in the inpatient setting. Table 1 describes the top drug classes that were overridden. The override rate for monoclonal antibodies was 98.1% (n=51) and included for comparison with the top drug classes in Table 1. The number of alerts overridden in the drug class 'Opioids' was 87.2% (n=58,348) followed by the class 'Non-antibiotic sulfonamides (without coxibs)' at 84.4% (n=514), and 'Non-antibiotic sulfonamides (coxibs)' at 81.6% (n=129). The override rate for 'Sulfonamide antibiotics' was far lower than other sulfonamides (59.6%, n=627). The 'Other drugs' category contained individual drugs that were collectively overridden 74.5% (n=24,463); drug allergies triggered for the top five 'other drugs' included paracetamol (4.4%, n=4,375), metoprolol (1.6%, n=2,168), prochlorperazine (1.1%, n=1,444), simvastatin (0.9%, n=1,154), and rosuvastatin (0.6%, n=826). The drug class 'Opioids' generated the highest number of total orders in the study period and were the second highest class to trigger an alert; orders for non-antibiotic sulphonamides triggered the highest percentage of alerts.

Table 1. Top drug classes that were overridden in the inpatient setting

| Drug class | Total No. Orders | Alerts Triggered | | Percentage of orders that triggered an alert | Alerts Overridden | |
|---|------------------|------------------|--------------------------------------|--|-------------------|------|
| | | n | % (no. alerts / total no. alerts) | | n | % |
| Monoclonal antibodies | 1,276 | 52 | 0 | 4.1 | 51 | 98.1 |
| Opioids | 428,074 | 66,949 | 50.9 | 15.6 | 58,348 | 87.2 |
| Non-antibiotic sulfonamides (without coxibs) | 152,286 | 609 | 0.5 | 0.4 | 514 | 84.4 |
| Non-antibiotic sulfonamides (coxibs) | 624 | 158 | 0.1 | 25.3 | 129 | 81.6 |
| Cephalosporins & other beta-lactams | 80,615 | 17,978 | 13.7 | 22.3 | 14,543 | 80.9 |
| Aspirin & NSAIDs* (without coxibs) | 139,069 | 8,730 | 6.6 | 6.3 | 6,809 | 78 |
| ACE inhibitorsξ & angiotensin receptor blockers | 46,319 | 724 | 0.6 | 1.6 | 560 | 77.3 |
| Penicillins | 26,999 | 2,379 | 1.8 | 8.8 | 1,679 | 70.6 |
| Sulfonamide antibiotics | 12,536 | 1,052 | 0.8 | 8.4 | 627 | 59.6 |
| Contrast media | 10,984 | 161 | 0.1 | 1.5 | 89 | 55.3 |
| Other drugs | 4,471,594 | 32,823 | 24.9 | 0.7 | 24,463 | 74.5 |
| Total | 5,370,376 | 131,615 | 100 | 2.45 | 107,723 | 81.9 |

* = Non-steroidal anti-inflammatory drugs (NSAIDs)

ξ = Angiotensin-converting-enzyme (ACE) inhibitors

In the outpatient setting, the number of drug allergy alerts displayed was only a fifth of those displayed over the same time period in the inpatient setting (n=26,408). The providers indicated that the *'patient did not have this allergy, will discontinue pre-existing allergy'* in over 5% (n=1,418) of cases. Overall alert override rate was statistically lower in the outpatient setting (77%, n=20,345), $p<.0001$. Table 2 describes the top drug classes that were overridden in the outpatient setting. There was only one alert for monoclonal antibodies, which was not overridden. The alert override rate for 'Non-antibiotic sulfonamides (coxibs)' (87.2%, n=157) was higher than for 'Non-antibiotic sulfonamides (without coxibs)' (81.9%, n=521), which differed from the inpatient setting. The top five drugs triggered the most frequently in the 'Other drugs' category were pravastatin (4%, n=1,046), guaifenesin (3.8%, n=998), rosuvastatin (3.7%, n=983), azithromycin (3.1%, n=827), and simvastatin (3.1%, n=821); all drugs in this category were collectively overridden 79.3% (n=11,092). Similar to the inpatient setting, the drug class 'Opioids' also generated the highest number of total orders in the study period but triggered a lower number of alerts per percentage of orders (5.1% compared to 15.64% in the inpatient setting). The drug class 'Non-antibiotic Sulphonamides' was also found to trigger the highest number of alerts per percentage of orders (10.67 %) in the outpatient setting. The percentage of orders for the drug class 'Cephalosporins & other beta-lactams' triggered a much higher number of alerts in the inpatient setting (22.3%) than the outpatient setting (8.8%).

Table 2. Top drug classes that were overridden in the outpatient setting

| Drug class | Total No. Orders | Alerts Triggered | | Percentage of orders that triggered an alert | Alerts Overridden (n=107,812) | |
|---|------------------|------------------|--------------------------------------|--|-------------------------------|------|
| | | n | % (no. alerts / total no. alerts) | | n | % |
| Non-antibiotic sulfonamides (coxibs) | 1,687 | 180 | 0.7 | 10.7 | 157 | 87.2 |
| Opioids | 106,972 | 5,451 | 20.6 | 5.1 | 4,605 | 84.5 |
| Non-antibiotic sulfonamides (without coxibs) | 78,947 | 636 | 2.4 | 0.8 | 521 | 81.9 |
| Aspirin & NSAIDs* (without coxibs) | 89,590 | 2,443 | 9.3 | 2.7 | 1,827 | 74.8 |
| ACE inhibitorsξ & angiotensin receptor blockers | 76,639 | 597 | 2.3 | 0.8 | 438 | 73.4 |
| Cephalosporins & other beta-lactams | 15,933 | 1,400 | 5.3 | 8.8 | 1,009 | 72.1 |
| Penicillins | 50,351 | 1,365 | 5.2 | 2.7 | 611 | 44.8 |
| Sulfonamide antibiotics | 24,412 | 347 | 1.3 | 1.4 | 85 | 24.5 |
| Other drugs | 1,550,659 | 13,988 | 53 | 0.9 | 11,092 | 79.3 |
| Monoclonal antibodies | 340 | 1 | 0 | 0.3 | 0 | 0 |
| Contrast media | 1 | 0 | 0 | 0 | 0 | 0 |
| Total | 1,995,531 | 26,408 | 100.1 (rounded to 100) | 1.3 | 20,345 | 77.0 |

* = Non-steroidal anti-inflammatory drugs (NSAIDs)

ξ = Angiotensin-converting-enzyme (ACE) inhibitors

Reasons for alert overrides

In the inpatient setting, the most common free-text reason for overriding drug allergy alerts was '*patient has taken previously without allergic reaction*' (Category 2, 57.4%, n=61,858), followed by '*physician aware*' (Category 7, 17.2%, n=18,583), and '*low risk cross sensitivity, will monitor*' (Category 3, 12.3%, n=13,202) (Table 3). In 1,386 (1.3%) of cases, the provider commented that they were administering the drug as per the desensitization protocol (Category 17, 0.9%, n=927) or that they were "*testing*" or "*trialing*" the drug on the patient (Category 19, 0.4%, n=459). In 224 (0.2%) of cases, the provider wrote that they would "*administer slowly*" (Category 11, 0.1%, n=112), "*discontinue the drug*" (Category 12, 0.05%, n=57), "*will discontinue if the patient experiences an adverse reaction*" (Category 13, 0.03%, n=37), or "*adjust dose*" (Category 9, 0.02%, n=18). According to providers, the patient either explicitly requested the drug or agreed to the drug being prescribed in 46 (Category 28, 0.04%) of cases.

Similarly in the outpatient setting, the most common coded reason given was '*patient has taken previously without allergic reaction*' (56.9%, n=11,594), followed by '*low risk cross sensitivity, will monitor*' (27.9%, n=5,685), and '*no reasonable alternatives*' (4.9%, n=986). Providers chose the coded reason '*other*' in 10% (n=2,037) of cases (Table 3). In the outpatient setting, a far lower number of providers commented "*physician aware*" (0.11%, n=23), that they were administering the drug as per the desensitization protocol (0.08%, n=17) or that they were "*testing*" or "*trialing*" the drug on the patient (0.07%, n=14). Almost the same percentage of providers in the outpatients setting as in the inpatient setting stated that they would "*discontinue the drug*" (0.04%, n=8) or "*discontinue if the patient experiences an adverse reaction*" (0.03%, n=7).

Table 3. Reasons given by providers for overriding drug allergy alerts

| Override reasons given by provider | Inpatient Category | Inpatients N (%) | Outpatients N (%) |
|---|-----------------------|---------------------|----------------------|
| Patient has taken previously without allergic reaction / patient has tolerated previously | 2 | 61,858 (57.4) | 11,594 (56.9) |
| Physician aware | 7 | 18,583 (17.2) | 23 (0.1) |
| Low risk cross sensitivity, will monitor | 3 | 13,202 (12.3) | 5,685 (27.9) |
| OK | 15 | 3,421 (3.17) | 20 (0.1) |
| No reasonable alternative / needed | 5 | 1,664 (1.5) | 986 (4.9) |
| Other (free-text reason provided) | 30 | 9,084 (8.4) | 2,037 (10) |
| Total | | 107,812 | 20,345 |

Appropriateness of alert overrides

In the inpatient setting, 96.5% (193/200) of the chart-reviewed alert overrides were judged to be appropriate. Of the 193 alerts that were overridden appropriately, three displayed the reaction ‘anaphylaxis’. Three of the seven specific alert overrides that were considered by the two independent reviewers to have been possibly inappropriately overridden were discussed with a third reviewer (D.W.B) before reaching final agreement; an inadequate explanation was provided in the free-text field for one of these overrides, and reviewers were unable to verify on chart review whether the patient received the drug and had an allergic reaction for a second of these overrides. In the outpatient setting, 94% (47/50) of alert overrides were considered appropriate. Three specific alert overrides were discussed with a third reviewer (D.L.S) before reaching final agreement (that they were inappropriate); in each of these cases, the provider chose the reason ‘other’ and provided no free-text explanation; the third reviewer was unable to verify on chart review whether the

patient had received the drug already without having an allergic reaction.

Types of reactions

The most common type of reaction displayed in the alert warnings in the inpatient setting related to the skin (27.4% of the total), which included rash (14.6%, n=19,172), hives (7.1%, n=9,353), and itching (5.7%, n=7,547). Over 21% of alerts (n=27,650) indicated some kind of gastro-intestinal upset and were overridden in 85.3% (n=23,599) of the cases. **Table 4** describes the top 10 reactions that were displayed to providers and the override rates of these specific types of potential reactions. The percentages shown in Columns 2 and 4 show the fraction of how many alerts warned against the risk of the respective reaction in the Inpatient and Outpatient Settings, respectively. The percentages shown in the Columns 3 and 5 present the fraction of how many of the respective alerts were overridden by the providers in the Inpatient and Outpatient Settings, respectively. The number of alerts that displayed '*shortness of breath*' and '*anaphylaxis*' in the Inpatient Setting was 2,541 (1.9%) and 2,373 (1.8%) respectively, and these were overridden in 81.2% (n=2,063) and 70.9% (n=1,682) of cases.

In the outpatient setting, over 22.8% of alerts (n=6,009) displayed gastro-intestinal upset and 81.9% (n=4,902) of these were overridden. Skin reactions were also commonly displayed (19.6%, n=5,167); these included rash (11.2%, n=2,956), hives (5.1%, n=1,353), and itching (3.2%, n=858), and were overridden on average 66% (n=3,413). The number of alerts that displayed '*shortness of breath*' and '*anaphylaxis*' was far lower in the outpatient setting and overridden much less 70.8% (n=289) and 56% (n=130), respectively.

Table 4. Top 10 reactions displayed to providers and the override rates of these alerts

| Reactions | Inpatient alerts N (%) | Inpatient overrides N (%) | Outpatient alerts N (%) | Outpatient overrides N (%) |
|----------------------------|------------------------------|---------------------------------|-------------------------------|----------------------------------|
| Unknown | 29,125 (22.1) | 23,745 (81.5) | 3,031 (11.5) | 2,038 (67.2) |
| Gastro-intestinal upset | 27,650 (21) | 23,599 (85.3) | 6,009 (22.8) | 4,902 (81.6) |
| Rash | 19,172 (14.6) | 15,659 (81.7) | 2,956 (11.2) | 1,868 (63.2) |
| Hives | 9,353 (7.1) | 7,539 (80.6) | 1,353 (5.1) | 906 (67) |
| Itching | 7,547 (5.7) | 6,430 (85.2) | 858 (3.2) | 639 (74.5) |
| Mental status change | 6,685 (5.1) | 5,610 (83.9) | 924 (3.5) | 765 (82.8) |
| Angioedema | 4,173 (3.2) | 3,136 (75.1) | 913 (3.5) | 613 (67.1) |
| Shortness of breath | 2,541 (1.9) | 2,063 (81.2) | 408 (1.5) | 289 (70.8) |
| Anaphylaxis | 2,373 (1.8) | 1,682 (70.9) | 232 (0.9) | 130 (56) |
| Myalgia | 2,195 (1.7) | 1,891 (86.2) | 2,224 (8.4) | 1,998 (89.8) |
| Total | 110,814 (84.2) | 91,354 (82.4) | 18,908 (71.6) | 14,148 (74.8) |

DISCUSSION

Over the last 20 years, there has been an increase in the drug allergy alert override rate at this hospital only from about 50% in 1995,¹⁶ to 81.9% (inpatients) and 77% (outpatients) found in this study. These high override rates are similar to those of previous studies which analyzed orders from the same and different hospitals,¹⁶⁻¹⁸ although a little lower than the 91% override rate found by Weingart et al. in the outpatient setting.¹³ Overall, these data suggest that better documentation and updating of allergy records are needed in patient's health records to ensure that these are correct and up-to-date, and further studies are needed to improve the sensitivity and specificity of allergy alerts, thus helping to specify which alerts should be turned on and which should be turned off.

A few classes accounted for a large proportion of this high drug allergy override rate, suggesting it could be readily addressed. Opiates in particular were one of the most frequently ordered and overridden drug classes, similar to other studies.^{11,17} A random sample of alert overrides in both settings were reviewed and over 94% were considered appropriate. Cross-reactivity in particular is an issue; many patients have a reaction to one opiate but it is not clear that providers need warnings when others are given. Hsieh et al also reviewed the charts of a random subset of patients and found that none of the adverse drug events attributed to the overridden drug alert were preventable.¹⁷ The authors concluded that clinical need for the drug outweighed the risk of a serious allergic reaction, thus the drug allergy alert overrides were deemed clinically justifiable.

Providers must ensure that patient self-reported allergies or intolerances are documented correctly in the electronic patient record and frequently updated. Many different types of providers can usually enter this information into the system at the time of the patient's visit.¹³ Hsieh et al. found that the degree of completeness of patients' allergy lists was generally poor in the CPOE system.¹⁷ The accuracy of the information included in these lists was also questionable, and it was often unclear whether the specific reactions were IgE or non-IgE-mediated. Often, opioid side effects are recorded as allergies.¹¹ This is because the reporting of allergies and intolerance and adverse effects are handled through the same mechanism. This may be acceptable, but applications should ask providers to distinguish between them when they are being reported. Hsieh et al also found that nausea and vomiting, which are not true IgE-mediated reactions, were the most common adverse drug reactions documented to narcotics in the patient's allergy list.¹⁷ Further, in our study, the provider indicated that the patient (for whom a drug-allergy alert was triggered) did

not actually have the allergy in 4,020 (2.5%) of cases. Providers overrode more than half of all drug allergy alerts giving the reason that the *'patient has taken previously without allergic reaction,'* with the scenario most often being that the patient had an allergy recorded to a drug but the patient was taking it without difficulty. Similar findings were reported in another study conducted in the inpatient setting with providers choosing the same reason (49%) when overriding drug allergy alerts.¹⁹ On selecting this override reason, it should be possible to encourage providers to deactivate allergies by providing an automatic link for providers to update the drug allergy list. Additional studies are also needed to look at the more efficient knowledge management of the drug allergy databases.

One of the most common override reasons given in both settings was *'low risk cross sensitivity, will monitor'*. Previous studies have revealed how the majority of drug allergy overrides (90%) occur when the allergen and drug belong to the same family but were not identical (e.g., codeine and hydromorphone).¹⁷ For some of the non-exact drug allergy matches presented in study by Hsieh et al., the likelihood of an adverse reaction occurring to the patient on administration of the drug was felt to be extremely low. For example, the drug classes *'non-antibiotic sulfonamides'* were overridden over 81% of the time in both inpatient and outpatient settings. Strom et al. conducted a retrospective cohort study examining patients who previously had an allergic reaction to a sulfonamide antibiotic and found that less than 10% of these patients had an allergic reaction to a sulfonamide non-antibiotic (e.g., celecoxib).²⁰ The authors concluded that this association appeared to be due to a predisposition to allergic reactions in general rather than to cross-reactivity with sulfonamide-based drugs. In other words, these patients were no more likely to have a subsequent hypersensitivity reaction to a drug containing a sulfonamide functional group (e.g.,

furosemide) than to a drug that did not (e.g., penicillin).²⁰ This highlights the importance of incorporating clinical evidence on drug cross-reactivity into the decision support system, rather than relying on pharmacologic or structural similarity.²¹

Over 70% of the drug allergy alerts that displayed the reaction ‘anaphylaxis’ in this study were overridden. This is a worrying finding, as a true allergic reaction severe enough to cause anaphylaxis will most often occur within minutes or hours of exposure. It is unclear from our study whether any of these patients (for whom an alert was triggered) had previously experienced such a reaction, or went on to experience any adverse events as a result of taking the prescribed drug (beyond the scope of this study). Identification of patients with a previous history of anaphylaxis is important, as over one third of these patients could suffer a serious recurrence.²² However, with recent figures suggesting that the rate of anaphylaxis hospitalizations is on the rise in both the U.K. and U.S.,^{23,24} further research is needed to explore why providers chose to override these alerts. In the inpatient setting, Topaz et al reported how only about one-tenth of the alerts for potentially life threatening reactions (e.g., anaphylaxis) were based on an exact match between the allergy and prescribed drug, while the rest were based on either the cross-sensitivity or allergy group.¹⁴ These findings further suggest that patients’ medical records may contain inaccurate allergy information or the clinical evidence on drug cross-reactivity may need to be updated. This may be of particular importance to pediatric patient populations whose drug allergy prevalence varies substantially from adults.²⁵

This study was undertaken within a single healthcare delivery system using two different homegrown prescribing systems (BICS and LMR) and, as such, is difficult to assess how generalizable the results are to other sites. However, our

proportions are comparable with the results from different sites published by other research groups. The random sample of alert overrides reviewed for appropriateness was small compared with the total number of alerts triggered, and the accuracy of this review was reliant on the completeness of information contained in the patients' charts. We did not control for the possibility that some alerts may have been repeated for the same patient. This paper presents the reasons why providers chose to override drug allergy alerts. Although providers in the inpatient setting enter this reason in the free-text field provided, providers in the outpatient setting selected one of the coded reasons provided. Notwithstanding these limitations, we believe that our findings have important implications for allergy alerting in other systems.

CONCLUSION

Drug allergy alerts are one of the most frequently displayed alerts in CPOE systems. There is little evidence, however, regarding the efficacy of these alerts. Most of the drug allergy alerts in this study were overridden, with over 94% of a sub-sample of alert overrides considered appropriate. Some of the rules for these alerts should be carefully reviewed and modified, or removed. It is very important that the information contained in patients' drug allergy lists and clinicians encouraged to deactivate allergies by providing them with an automatic link to update their patients' drug allergy lists. The clinical evidence on drug cross-reactivity in the decision support systems should be updated. Finally, we found that providers overrode the majority of drug allergy alerts, including those that warned against the risk of 'anaphylaxis'. More needs to be done to understand providers' reasoning in overriding such alerts and reduce the possible threat to patient safety.

REFERENCES

1. American College of Allergy Asthma and Immunology. Allergy Facts. Available from: <http://acaai.org/news/facts-statistics/allergies> (Accessed on 2nd January 2016).
2. World Allergy Organization. *White Book on Allergy 2011-2012. Executive Summary*. By Prof. Ruby Pawankar, MD, PhD, Prof. Giorgio Walkter Canonica, MD, Prof. Stephen T. Holgate, BSc, MD, DSc, FMed Sci and Prof. Richard F. Lockey, MD. World Allergy Organization. 2011
3. Balas EA, Weingarten S, Garb CT, Blumenthal D, Boren SA, Brown GD. Improving preventive care by prompting physicians. *Arch Intern Med*. 2000;160(3):301-308.
4. Bates DW, Gawande AA. Patient safety: Improving safety with information technology. *New Engl J Med*. 2003;348(25):2526-2534.
5. Bates DW, Leape LL, Cullen DJ, et al. Effect of computerized physician order entry and a team intervention on prevention of serious medication errors. *J Am Med Assoc*. 1998;280(15):1311-1316.
6. Kaushal R, Shojania KG, Bates DW. Effects of computerized physician order entry and clinical decision support systems on medication safety: a systematic review. *Arch Intern Med*. 2003;163(12):1409-1416.
7. Kaushal R, Jha AK, Franz C, et al. Return on investment for a computerized physician order entry system. *J Am Med Inform Assn*. 2006;13(3):261-266.
8. Slight SP, Seger DL, Nanji KC, et al. Are we heeding the warning signs? Examining providers' overrides of computerized drug-drug interaction alerts in primary care. *PLoS One*. 2013;8(12):e85071.
9. Nanji KC, Slight SP, Seger DL, et al. Overrides of medication-related clinical decision support alerts in outpatients. *J Am Med Inform Assoc*. 2014;21(3):487-491.
10. Smithburger PL, Buckley MS, Bejian S, Burenheide K, Kane-Gill SL. A critical evaluation of clinical decision support for the detection of drug-drug interactions. *Expert opinion on drug safety*. 2011;10(6):871-882.
11. Ariosto D. Factors Contributing to CPOE Opiate Allergy Alert Overrides. *AMIA Annu Symp Proc*. 2014;2014:256-265.

12. Drug Allergies. Thong B, Vervloet D (Updated 2014). World Allergy Organization website. . Available from:
http://www.worldallergy.org/professional/allergic_diseases_center/drugallergy/
[Accessed on 2nd January 2016.
13. Weingart SN, Toth M, Sands DZ, Aronson MD, Davis RB, Phillips RS. Physicians' decisions to override computerized drug alerts in primary care. *Arch Intern Med.* 2003;163(21):2625-2631.
14. Topaz M, Seger, D., Slight, S.P., Goss, F., Lai, K., Wickner, P.G., et al. Rising drug allergy alert overrides in electronic health records: an observational retrospective study of a decade of experience. *J Am Med Inform Assoc.* 2015 Nov 17. pii: ocv143. doi: 10.1093/jamia/ocv143
15. James CW, Gurk-Turner C. Cross-reactivity of beta-lactam antibiotics. *Proc (Bayl Univ Med Cent).* 2001;14(1):106-107.
16. Abookire SA, Teich JM, Sandige H, et al. Improving allergy alerting in a computerized physician order entry system. *Proc AMIA Symp.* 2000:2-6.
17. Hsieh TC, Kuperman GJ, Jaggi T, et al. Characteristics and consequences of drug allergy alert overrides in a computerized physician order entry system. *J Am Med Inform Assoc.* 2004;11(6):482-491.
18. Lin CP, Payne TH, Nichol WP, Hoey PJ, Anderson CL, Gennari JH. Evaluating clinical decision support systems: monitoring CPOE order check override rates in the Department of Veterans Affairs' Computerized Patient Record System. *J Am Med Inform Assoc.* 2008;15(5):620-626.
19. Huntman L, Ward L, Read D, Jolly M, Heckman M. Analysis of allergy alerts within a computerized prescriber-order-entry system. *Am J Health Syst Pharm.* 2009;66(4):373-377.
20. Strom BL, Schinnar R, Apter AJ, et al. Absence of cross-reactivity between sulfonamide antibiotics and sulfonamide nonantibiotics. *N Engl J Med.* 2003;349(17):1628-1635.
21. Carspecken CW, Sharek PJ, Longhurst C, Pageler NM. A clinical case of electronic health record drug alert fatigue: consequences for patient outcome. *Pediatrics.* 2013;131(6):e1970-1973.
22. Mullins RJ. Anaphylaxis: risk factors for recurrence. *Clin Exp Allergy.* 2003;33(8):1033-1040.

23. Ma L, Danoff TM, Borish L. Case fatality and population mortality associated with anaphylaxis in the United States. *J Allergy Clin Immunol*. 2014;133(4):1075-1083.
24. Turner PJ, Gowland MH, Sharma V, et al. Increase in anaphylaxis-related hospitalizations but no increase in fatalities: an analysis of United Kingdom national anaphylaxis data, 1992-2012. *J Allergy Clin Immunol*. 2015;135(4):956-963 e951.
25. Thong BY, Tan TC. Epidemiology and risk factors for drug allergy. *Br J Clin Pharmacol*. 2011;71(5):684-700.